

國立中正大學八十七學年度碩士班考試試題

所 別：化學工程學系

科 目：工程數學

1. An iterative equation is described by the equation $\mathbf{x}_{n+1} = \mathbf{A}\mathbf{x}_n$,

where $\mathbf{A} = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix}$. If $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, compute \mathbf{x}_{11} . (10 points)

2. (a) Let $f(t)$ have Fourier transform $\phi(\omega)$. Show that

$$\int_{-\infty}^{\infty} |\phi(\omega)|^2 d\omega = 2\pi \int_{-\infty}^{\infty} |f(t)|^2 dt \quad (6 \text{ points})$$

(b) Use Laplace transform to solve $\frac{d^2x}{dt^2} + \frac{dx}{dt} + 2x = e^{3t}$ with initial conditions $x = 1$ and $\frac{dx}{dt} = 1$ for $t = 0$. (4 points)

3. (a) Briefly state the difference between the Laplace and the Fourier Transforms. (5 points)

(b) Briefly state one solution method for a non-homogeneous set of ordinary differential equations $\frac{d\mathbf{x}}{dt} = \mathbf{A}\mathbf{x} + \mathbf{f}(t)$, where $\mathbf{x} \in \mathbb{R}^n$.

(5 points)

(c) For 4-dimensional vectors \mathbf{a} , \mathbf{b} , \mathbf{c} , and \mathbf{d} , how to determine whether they can form a basis in the 4-dimensional space.

(5 points)


4. A boat A moves along the y axis with constant speed a . A second boat B which moves in the left-hand half of the x - y plane with constant speed b and always points directly at boat A .

(a) Show that the path of boat B is described by the differential equation

$$x \frac{d^2y}{dx^2} = -\frac{a}{b} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \quad (9 \text{ points})$$

(b) Solve the path of boat B .

(6 points)



國立中正大學八十七學年度碩士班考試試題

所 別：化學工程學系

科 目：工程數學

5. The time evolution of velocity profile for the Couette flow is described by the partial differential equation,

$$\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial y^2}, \quad (\nu \text{ is a constant})$$

subject to the following initial and boundary conditions:

$$\begin{aligned} u(y, 0) &= 0, & 0 < y < 1, \\ u(0, t) &= U, & u(1, t) = 0, & t > 0. \end{aligned}$$

- (i) Solve the steady-state velocity profile $u(y)$ (5 points).
(ii) Solve the transient velocity profile $u(t, y)$ (15 points). (Hint: You may use the result in (i) and a change of variable to make the boundary conditions homogeneous)

6. Here we are interested in finding approximate solutions for a first-order, nonlinear, ordinary differential equation,

$$x \frac{dx}{dt} - (1+t)x = \cos(2\pi t), \quad x(0) = 0, \quad x(1) = 1.$$

- (i) First show that (5 points)

$$\left. \frac{dx}{dt} \right|_x = \frac{x(t+\Delta t) - x(t-\Delta t)}{2\Delta t} + O[(\Delta t)^2].$$

- (ii) Use the above relation to find an approximate solution for $x(0.5)$ (10 points).

7. Verify the Gauss-Ostrogradskii divergence theorem,

$$\int_V \nabla \cdot \mathbf{u} \, dV = \int_S \mathbf{n} \cdot \mathbf{u} \, dS, \quad (\mathbf{n} \text{ is the outwardly directed normal vector of } dS)$$

for $\mathbf{u} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ in the region bounded by $x = \pm 1, y = \pm 1, z = \pm 1$ (15 points).

第二頁
(共二頁)